




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C11D 3/39, C07D 227/087</b>		A1	(11) International Publication Number: <b>WO 95/14759</b>
			(43) International Publication Date: 1 June 1995 (01.06.95)
(21) International Application Number: <b>PCT/GB94/02583</b>		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).	
(22) International Filing Date: 25 November 1994 (25.11.94)			
(30) Priority Data:			
9324261.8	25 November 1993 (25.11.93)	GB	
9325777.2	16 December 1993 (16.12.93)	GB	
9326382.0	23 December 1993 (23.12.93)	GB	
9326385.3	23 December 1993 (23.12.93)	GB	
9326383.8	23 December 1993 (23.12.93)	GB	
9326384.6	23 December 1993 (23.12.93)	GB	
(71) Applicant (for all designated States except US): <b>WARWICK INTERNATIONAL GROUP LIMITED [GB/GB]; Wortley Moor Road, Leeds LS12 4JE (GB).</b>		<b>Published</b> <i>With international search report.          Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(72) Inventors; and			
(75) Inventors/Applicants (for US only): <b>CROWTHER, Jan, Darrel [GB/GB]; Vaughn Close, Prestatyn, Clwyd LL19 8SH (GB). HAQUE, Ekram [GB/GB]; 4 Ash View, Aston Park, Hawarden, Deeside, Clwyd CH5 1BL (GB). COLES, Barrie [GB/GB]; Montrose, Pant Asaph, Holywell, Clwyd CH8 8PN (GB).</b>			
(74) Agent: <b>GILL JENNINGS &amp; EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).</b>			

(54) Title: **BLEACHING COMPOSITIONS**

## (57) Abstract

Bleach activators and the peroxygen-based oxidising species formed in situ from a peroxygen source and a bleach activator, the bleach activator having formula (I) in which R represents an optionally substituted alkenyl function having at least 2 carbon atoms in the alkenyl chain which forms a cyclic structure with -N- and -C(O)- and R<sup>1</sup> represents a C<sub>1-30</sub> alkyl or aryl group; or has the formula -R<sup>2</sup>-COX where X may be OH, OM (where m is a metal ion), -ONR<sup>3</sup><sub>4</sub> (where R<sup>3</sup> are each independently selected from H and C<sub>1-4</sub> alkyl), a halogen atom, amine, C<sub>1-30</sub> alkyl or alkoxy or a leaving group and R<sup>2</sup> is a C<sub>1-30</sub> alkylene or arylene group with the proviso that where R is (CH<sub>2</sub>)<sub>5</sub>, R<sup>1</sup> represents a C<sub>3-9</sub> branched alkyl group or a C<sub>10-30</sub> alkyl or aryl group or a C<sub>9-10</sub> branched aralkyl group; or has the formula -R<sup>2</sup>-COX and R<sup>2</sup> is a C<sub>3-10</sub> branched alkyl group or C<sub>11-30</sub> alkylene or arylene group, or a C<sub>9-10</sub> branched alkarylene. Compositions including these bleach activators in combination with conventional bleach activators such as tetraacylated alkylene diamines, particularly TAED or SNOBS give particularly useful cleaning properties.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

### Bleaching Compositions

This invention relates to the in situ production of peroxygen-based oxidising species from a peroxygen source and an activator followed by the use of the product an oxidising agent, for instance in a detergent or as a bleach or a biocide.

It is known that percarboxylic acid compounds are useful as bleaching agents, general oxidising agents and biocides. Such compounds may be incorporated into for instance laundry detergents, hard surface cleaners, paper and pulp bleaching compositions, fabric and fibre bleaching compositions and even depilatory creams as well as biocidal compositions, for use in agriculture.

It is also known to use a combination of peroxygen bleach precursor (or peroxygen source) and bleach activator in the same or separate compositions to form the oxidising species. The bleach activators are acyl-releasing agents. Suitable bleach activators are O-acyl or N-acyl compounds.

However, there is still a need for further bleach activators which will provide good bleaching and may also provide improved cleaning in detergent compositions.

In GB 855735 bleaching processes and compositions for textile material are described in which the bleaching composition is an aqueous solution comprising hydrogen peroxide and a reactive acyl organo-amide. A large range of compounds is included in this definition, including N-acyl lactams, of which the only exemplified lactam is N-acetyl caprolactam.

GB 1596313 discloses detergents for laundering textiles in which the compositions contain bleaching agents and an activator. The only activator mentioned is N-acetyl caprolactam. N-acetyl caprolactam is also mentioned in GB 2189520 A which relates to washing and bleaching processes. A typical example of a bleach precursor is given as N-acetyl caprolactam. Caprolactam in its N-acetylated form

is also known as an ingredient in detergent compositions from SU 1664826.

In EP-A-0122763, bleach compositions are described in which an activator is adsorbed onto sodium perborate monohydrate. The activators claimed are any O-acyl or N-acyl activator and a large range of suitable compounds are disclosed. Within these compounds, N-acyl caprolactam is mentioned, where the acyl group is  $R^aCO$  in which  $R^a$  represents hydrogen or an alkyl group of chain length  $C_1$  to  $C_9$  aliphatic or aromatic group, optionally substituted by an alkyl or carboxylic acid group or the acyl is a di-acyl derived from a dicarboxylic acid having from 4 to 12 carbon atoms.


The present invention relates to novel bleach activators which provide improved cleaning properties in detergent compositions.

In accordance with the present invention there is provided a bleach activator comprising an optionally substituted N-acyl heterocycle having the formula I:




in which R represents an optionally substituted alkenyl function having at least 2 carbon atoms in the alkenyl chain which forms a cyclic structure with -N- and -C(O)- and  $R^1$  represents a  $C_{1-30}$  alkyl or aryl group; or has the formula  $-R^2-COX$  where X may be OH, OM (where M is a metal ion),  $-ONR^3$ , (where  $R^3$  are each independently selected from H and  $C_{1-4}$  alkyl), a halogen atom, amine,  $C_{1-30}$  alkyl or alkoxy or a leaving group and  $R^2$  is a  $C_{1-30}$  alkylene or arylene group with the proviso that where R is  $(CH_2)_5$ ,  $R^1$  represents a  $C_{3-9}$  branched alkyl group or a  $C_{10-30}$  alkyl or aryl group or a  $C_{9-10}$  branched aralkyl group; or has the formula  $-R^2-COX$  and  $R^2$  is a  $C_{3-10}$  branched alkyl group or  $C_{11-30}$  alkylene or arylene group, or a  $C_{9-10}$  branched alkarylene.


Thus, the compounds of the formulae I may be optionally substituted N-acyl 3-membered heterocycles (2-azetidinones) of the formula II:




N-acyl 5-membered heterocycles, N-acyl pyrrolidones  
10 having the formula (III):




they may be N-acyl 6-membered rings of the formula  
(IV):




N-acyl 7-membered rings caprolactams of the formula  
(V):



N-acyl 8-membered heterocycles (2-azocyclooctanones)  
of the formula (VI):



40 or N-acyl 9-membered heterocycles (2-azocyclononanones) of  
the formula VII:



(VII)

5

or N-acyl heterocycles having 10-membered rings or above. In each of the formulae listed above,  $\text{R}^1$  is as defined for formula I above.

10 Where the bleach activator is a 7-membered ring, preferably  $\text{R}^1$  is a  $\text{C}_{10-28}$ , or  $\text{C}_{12-26}$  alkyl group, most preferably  $\text{R}^1$  is a  $\text{C}_{14-24}$  alkyl group. Although it is not preferred,  $\text{R}^1$  may be a group linked to a substituent on the caprolactam ring to form a cyclic structure. Any branching  
15 may be with  $\text{C}_1$  or  $\text{C}_2$  groups or alternatively  $\text{C}_3$  or above alkyl groups. Where the bleach activator is other than a 7-membered ring, preferably  $\text{R}^1$  is a  $\text{C}_{2-14}$ , most preferably  $\text{C}_{7-12}$  group. Particularly preferred groups are aryl especially phenyl, which are optionally substituted, or alkyl groups,  
20 preferably straight chain, especially ethyl. Although it is not preferred,  $\text{R}^1$  may be a group linked to a substituent on the ring formed from  $-\text{R}-$ ,  $-\text{N}-$  and  $-\text{C}(\text{O})-$ , to form a cyclic structure.


Particularly preferred activators of the present  
25 invention for compounds of formula (II), (III), (IV), (VI) or (VII) are benzoyl N-substituted heterocycles. In the present invention, unless specified, the term alkyl(ene) includes alkynyl(ene) and alkenyl(ene) groups and these groups may be straight chain, branched or may include  
30 cycloalkyl, or groups. The alkyl groups may include heteroatoms such as N or O in the chain. As used herein, aryl(ene) includes phenyl(ene) and alkaryl(ene). The alkyl(ene) or aryl(ene) may include heterocyclic groups. They may be such substituted such as hydroxyl, lower  
35 alkyl (i.e. up to  $\text{C}_5$ ), amine, substituted amine (especially substituted with  $\text{C}_{1-30}$ , preferably  $\text{C}_{1-5}$  alkyl groups), acyl, acyloxy, alkoxy, aryl, aroyl, aryloxy, aroyloxy, halogen, amido, imido groups nitro, sulphonic

acid and the like they may be alkoxyated with ethers having C<sub>1-30</sub> alkyl groups, in particular, ethoxylated or propoxylated, or substituted as well as any other groups not adversely affecting the activity of the compound.


5 Where X is a leaving group, suitable leaving groups are for example any group which may be displaced from the bleach activator as a consequence of nucleophilic attack on the bleach activator. Generally leaving groups exert an electron attracting effect and are therefore weak bases.  
 10 Where X is a leaving group it preferably forms a compound the conjugate acid of which has a pKa in the range 4-13, preferably 7-11, most preferably 8-11. Good leaving groups are also those which form a stable entity to minimise the rate of back reaction. Suitable leaving groups include

15

20



and



and preferred leaving groups include








25

30

35


40

45




6

5



10



wherein  $\text{R}^5$  and  $\text{R}^6$  are hydrogen, or alkyl or aryl, preferably having from 1 to 14 carbon atoms;

$\text{R}^4$  is H or an alkyl chain containing from 1 to 8 carbon atoms;

$\text{R}^7$  is H, alkyl or aryl;

$\text{R}^9$  is alkyl or aryl, preferably having from 1-20, most preferably 1-14 carbon atoms; and

Y is H or a solubilising group.

The preferred solubilising groups in Y are  $-\text{SO}_3^-\text{M}_1^+$ ,  $-\text{COO}^-\text{M}_1^+$ ,  $-\text{SO}_4^-\text{M}_1^+$ ,  $(-\text{N}^+\text{R}^8)_2\text{X}_2^-$  and  $\text{O}=\text{N}(\text{R}^8)$

and most preferably  $-\text{SO}_3^-\text{M}_1^+$  and  $-\text{COO}^-\text{M}_1^+$  wherein  $\text{R}^8$  is an alkyl chain containing from 1 to 4 carbon atoms,  $\text{M}_1$  is a cation. Preferably,  $\text{M}_1$  is a metal ion as described above (M), or ammonium or a substituted ammonium cation, with sodium and potassium being most preferred, and  $\text{X}_2$  is a halide, hydroxide, methylsulfate or acetate anion.

On reaction of a bleach activator of the present invention with a peroxygen source, a more active bleaching species which may be a peracid is formed, in addition to unsubstituted heterocycle. Useful examples of bleach activators according to formula I include 5-membered ring heterocycles as defined in formula I where R is  $-(\text{CH}_2)_3$  and  $\text{R}^1$  is a phenyl, 4-methylphenyl, 4-nitrophenyl, 4-tertiary butylphenyl, 1-ethylphenyl, 3-pyridyl, pentyl, 4-chlorophenyl, 4-methoxyphenyl, octyl, phenoxyethyl, 2-nitrophenyl, 3-nitrophenyl or tertiary butyl group;

7-membered ring heterocycles as defined in formula I where R is  $-(\text{CH}_2)_5$  and  $\text{R}^1$  is 3,4 dicarboxyphenyl anhydride, 4-tertiarybutyl phenyl, tertiary butyl, 5-(N-



phthalimidyl)pentyl, tridecyl, phenoxymethyl, 3-(N,N-dimethylamino)phenyl, 4-(N,N-dimethylamino) phenyl, 1-ethylpentyl or heptadecyl;

8-membered ring heterocycles of formula I where R is  
5  $\{CH_2\}_6$  and R<sup>1</sup> is methyl or phenyl.

The present invention also includes a bleaching composition comprising one or mixtures of more than one of the bleach activators as defined above.

In the present invention the term "bleaching" is  
10 intended to include any process where the solution is used to remove or reduce unwanted colour in a substrate and/or reduce or remove non-coloured stains from fabric substrate and/or act as a disinfectant and the term "bleaching composition" is exemplified by but not limited to  
15 detergents, especially laundry detergents, dishwashing detergents, hard surface cleaners, including kitchen and bathroom cleaners, toilet cleaners, pipe cleaners, dry bleaches, compositions for bleaching wood, paper and pulp bleaching compositions, fabric and fibre bleaching  
20 compositions and oral hygiene compositions including mouthwashes, toothpastes and plaque-removing compositions and depilatory creams. In such bleaching compositions, there is included a precursor peroxygen source, that is a composition which releases perhydroxyl groups (OOH<sup>•</sup>).

25 The precursor peroxygen source may be hydrogen peroxide itself, but is alternatively an inorganic salt for instance a percarbonate or a perborate. Sodium or potassium salts are preferred, such as sodium perborate, or an organic peroxide such as benzoyl peroxide or urea  
30 peroxide.

Alternatively, bleaching compositions according to the invention may be used in conjunction with separate compositions which comprise such a bleach precursor.

The invention also includes a process in which a  
35 peroxygen source is reacted with one of the above activator compounds in aqueous solution to form a product solution

containing an oxidising product and the product solution is used as a bleach.

Where the peroxygen source is hydrogen peroxide itself, the concentration of hydrogen peroxide is preferably less than 70% (weight/volume) that is weight of hydrogen peroxide based on volume of water plus hydrogen peroxide plus other components in the mixture). Preferably the concentration is less than 60% weight by volume and more preferably less than 30% w/v. Where the product of the reaction is to be used in a domestic environment or other environment where it is difficult to take special precautions in handling the product, it is preferred for the concentration to be less than 15 or even less than 10% w/v. The concentration is usually at least 1% w/v more preferably at least 2% w/v.

Where the peroxygen source is other than hydrogen peroxide the concentration is preferably such as to give the equivalent available oxygen as the quoted concentrations of hydrogen peroxide. The concentration of peroxygen source in the aqueous liquid is for instance less than 10 M, preferably less than 5 M or sometimes even less than 3M down to 0.01 M. Preferably the concentration is at least 0.1M more preferably at least 0.2M. During the reaction, the temperature is preferably in the range 0 to 95°C, more preferably in the range 10 to 80°C, for instance in the range 20 to 60°C. The temperature in any subsequent oxidising step is preferably in the same ranges as the temperature during the reaction of the peroxygen source and the activator compound and is preferably substantially the same temperature specifically where the product solution is used for instance as a bleach or disinfectant. A particular advantage of using activators for the peroxygen source is that the oxidising product tends to be formed at a relatively low temperature, for instance 40°C or less, or even less than 30°C or 25°C which is advantageous for both safety and energy.

The compositions of the present invention are particularly useful in detergents, especially laundry detergents for both natural and synthetic fibres.

5 Generally, the reaction between the peroxygen source and the activator will take place under alkaline conditions, above pH 7 or 7.5. However, the reaction may also take place under acidic conditions. Generally the pH for the reaction will be from 6 to 14. Preferably it will be above 7.5 or even 8 and most preferably the pH will be  
10 from 9.5 to 12.5. Generally the bleaching compositions will be alkaline at a pH above 7.5. However, the composition may also contain a pH-modifier which may comprise an acid, a base and/or buffering material so that the desired pH can be obtained for the reaction between the  
15 precursor and the activator. For example, when the pH-modifier is an acidifying pH-modifier, it may comprise a polybasic organic acid such as a polybasic carboxylic acid, for example citric, succinic, adipic or sulphonic acid. Alternatively the pH-modifier may react with a by-product  
20 of the perhydrolysis reaction of the peroxygen source with the activator compound to produce an acid or base. Where perborate is used, borate is a by-product and so any component known to react with perborate to produce acidity for example a polyol, boric acid or sodium di-hydrogen  
25 phosphate can be used to provide the pH-modifier.

In the composition of the present invention, the peroxygen source and the activator compound may be provided in separate compositions. For instance, the peroxygen source may be in the form of an aqueous solution of  
30 peroxygen whilst the activator may be provided in the solid form. However, preferably the components are provided in a stable composition, which is preferably therefore substantially water free.

The composition may be in liquid form, for instance in  
35 a non-aqueous liquid medium, in which the components may be dissolved or dispersed. For instance particles of activator with protective coatings, for instance produced

by micro-encapsulation techniques or spray coating of solid activator, may be suspended in an aqueous, or non aqueous, solution of peroxygen source. As an alternative to a solution of peroxygen source that component may also be  
5 suspended in the liquid medium, either in a separate liquid phase or in particulate dispersed phase, particles of solid peroxygen source optionally being coated with a protective coating. Coated particles of either peroxygen source or activator may be disrupted or diluted into water or with  
10 abrasion.

Preferably the composition (or compositions) is in solid form, for instance as a mixture of particles of the individual components or, more preferably, comprising particles each of which comprise all of the components.  
15 Such particles may be provided by techniques the same or similar to those conventionally used in the laundry detergent industry, for instance including particles produced by spray drying liquid slurries, by granulation techniques using binders (for instance synthetic or natural  
20 polymers or derivatives) or by melt blending followed by extrusion or other techniques.

Preferably the composition contains the active ingredients in appropriate relative quantities so that when the composition is diluted (or the compositions are mixed)  
25 with water the first step of the reaction proceeds at the optimal rate and at the desired pH. The activator and peroxygen source are for instance present in relative amounts such that 5% to 150% of the stoichiometric amount of activator (for complete reaction with the peroxygen  
30 source) is provided. Preferably the amount of activator is 10 to 100%, more preferably 20 to 80% of the stoichiometric amount.

The new composition may include other additives, for instance stabilisers which stabilise the product before  
35 use, as well as stabilisers for the peroxy acid oxidising species formed in the reaction, especially heavy metal sequestrants. The new product may also include surfactants

to act as wetting agents and inorganic salts, for instance which affect the physical properties of the solid form or act as diluent. Other ingredients may be included depending on the final application of the reaction product, for instance perfumes, or agents to assist dissolution or dispersion of the product into water.

The N-substituted heterocyclic bleach activator of formula I may be simply admixed directly in with the other detergent composition compounds. Alternatively it could be coformulated with builders (for example phosphates, polyphosphates, sodium carbonate, silicates, zeolites clays, borax) complexing or sequestering agents (ethylenediamine tetraacetic acid, diethylenepentamine pentaacetic acid, nitrilotriacetic acid, citrates, phosphonates), enzymes, surfactants (linear alkylbenzene sulphonates, phenol sulphonates, alcohol ether sulphates, alcohol ethoxylates, amphoteric and cationic surfactants), soap, polyacrylates, hydrotropes, mineral oil and foam regulators.

The reaction product of the peroxygen source and the activator is preferably used immediately, without removal of any by-products or addition of other materials. Sometimes it may be desirable to add additional ingredients to the composition after the reaction between the peroxygen source and bleach activator such as pH-adjusters, wetting agents, additives to improve the action of the bleaching agent produced in the reaction, for instance co-disinfectants, biocides, abrasives etc.


The invention has been found to be surprisingly advantageous when the composition of the present invention comprises a peroxygen source and an N-substituted heterocycle of formula I as defined above, in combination with a conventional activator. The compositions have been found to exhibit enhanced bleaching effects. The compositions are especially effective as laundry detergents for both natural and synthetic fibres. Suitable conventional activators include any compound or mixture of

compounds already known as useful bleach activators. Particularly preferred conventional activators are tetraacylated alkylene diamines such as TAMD or TAED, tetraacylated glycol urils especially TAGU (tetraacetyl glycol uril), pentaacylated glucose, especially PAG (pentaacetyl glucose) and acylated (preferably C<sub>6-14</sub>) oxybenzene sulphonates, especially SNOBS (sodium nononoyl oxybenzene sulphonate).

It has been found that compositions incorporating both one or more of the N-acyl caprolactam activators of the present invention and a tetraacylated alkylene diamine, preferably TAED have particularly improved detergency properties.

In accordance with a further aspect of the present invention there is provided a bleaching composition comprising a peroxygen source, a conventional bleach activator, preferably a tetraacylated alkylene diamine and a bleach activator of the formula IX:

20



(IX)

in which R represents an alkenyl function having at least 2 carbon atoms in the alkenyl chain which forms a cyclic structure with -N- and -C(O)- and R<sup>1</sup> represents a C<sub>1-30</sub> alkyl or aryl group or has the formula -R<sup>2</sup>-COX where X is as defined for formula I above, and R<sup>2</sup> is a C<sub>1-30</sub> alkylene or arylene group. Preferably R<sup>1</sup> is a C<sub>7-30</sub> most preferably C<sub>7-15</sub> group, or has the formula -R<sup>2</sup>-COX where X is as defined above and R<sup>2</sup> is a C<sub>7-30</sub> group. Preferably R<sup>1</sup> or R<sup>2</sup> is straight chain alkyl(ene) or aryl(ene) group. Preferably R is a C<sub>5</sub> group or a C<sub>3</sub> group. Preferably R is an unsubstituted alkylene group, eg. -(CH<sub>2</sub>)<sub>5</sub> or (CH<sub>2</sub>)<sub>3</sub>.

A particularly preferred R<sup>1</sup> or R<sup>2</sup> group is an optionally substituted aryl(ene) group especially substituted phenyl(ene) group. A further preferred group includes bleach activators having the formula I above and

in which  $R^1$  represents a  $C_{10-30}$  or has the formula  $-R^2-COX$  where X is as defined above  $R^2$  is a  $C_{11-30}$  alkyl group and R comprises from 2 to 7  $-CH_2-$  groups. Without wishing to be bound by theory, it is postulated that the particular advantages which have been found for a combination of the N-acyl bleach activators of formula IX and I with acylated alkylene diamines, and in particular TAED, are due to the short chain nature of the peroxygen bleach compound produced using the alkylene diamine resulting in good hydrophilicity and the relatively long chain peroxygen bleach compounds formed from the activator of the present invention producing good hydrophobicity. Thus, the combination of the two activators used in combination gives good cleaning properties.

In a further embodiment of the invention, the conventional activator to be used in combination with the activator as defined in formula IX above comprises SNOBS (sodium nonanoyloxy benzene sulphate). In this embodiment of the invention, it is particularly preferred that the SNOBS is in combination with a bleach activator having the formula IX or I above, in which  $R^1$  represents a  $C_{1-9}$ , more preferably  $C_{2-6}$  alkyl group or has the formula  $-R^2-COX$  where X is as defined above and  $R^2$  is a  $C^{1-10}$  alkyl(ene) group, preferably a  $C_{2-6}$  alkyl group and R is a  $C_{2-7}$  group, preferably alkylene in which the  $CH_2$  groups of the lactam are unsubstituted. Preferably, the activator according to formula I given above is one in which R is a  $C_5$  group and  $R^1$  represents a  $C_{3-8}$  branched alkyl group.

Where the composition comprises at least two bleach activators, the first being a bleach activator of formula IX, preferably the activator of formula IX is present in an amount of at least 50% by weight total bleach activator, more preferably at least 70%, or even at least 80 or 90% by weight total bleach activator.

The invention is particularly directed to the manufacture of detergents for example used in washing

laundry and the compositions therefore preferably include conventional surfactants.

The novel bleach activators of the present invention can be prepared by any conventional organic synthesis route. They may be prepared by reacting the unsubstituted heterocycle with an acid chloride in which the acyl group is the desired acyl group for N-substitution on the unsubstituted heterocycle. The reaction is preferably carried out under alkaline conditions in a suitable organic solvent such as toluene. The reaction product can be crystallised or precipitated out of the solvent medium.

The invention is illustrated by way of the following examples:

Example 1

**Preparation of 4-nitrobenzoyl 2-pyrrolidone.**

A two litre flat-flanged flask equipped with thermometer, stirrer, dropping funnel and reflux condenser was charged with 292.0g of toluene, 2.78 mol of 2-pyrrolidone and 3.00 mol triethylamine. Separately, 2.78 mol 4-nitrobenzoyl chloride was dissolved in 400g of toluene and this mixture was added drop-wise to the contents of the reaction flask at room temperature. Once the addition was completed, the contents of the flask were refluxed for 4 hours. The reaction mixture was then cooled to room temperature and washed with  $2 \times 100\text{cm}^3$  water to remove the amine hydrochloride. The mixture was then filtered to give the first crop of product. A second crop of product may be obtained from the toluene layer if required by recrystallisation. The yield of 4-benzoyl 2-pyrrolidone was 474.2g (72.9%). The melting point of the product was 127-129°C and the purity as measured using HPLC was 98.7%. An NMR spectra of the product was obtained and the results were as follows:


- (1) doublet of doublets at around 8ppm (8.25 ppm and 7.70 ppm) indicative of 4-di-substituted aromatic hydrogen atoms ( $\text{H}_7$ - $\text{H}_{10}$ ).
- (2) quintuplet centred at around 2.17ppm ( $\text{H}_3$  and  $\text{H}_4$ ).



- (3) triplet centred at 3.98 ppm ( $H_1$  and  $H_2$ ).  
 (4) triplet centred at 2.62 ppm ( $H_5$  and  $H_6$ ).

5

10



15

An IR spectra of the product was obtained giving results: 2 bands at  $1746\text{cm}^{-1}$  and  $1666\text{cm}^{-1}$ , respectively, which are indicative of the two carbonyl bonds associated with cyclic imides.

#### Performance Testing

The bleach activator prepared in example 1 and several other bleach activators of the present invention as listed in Table 1, were tested for washing performance, comparisons being made using: (i) a blank incorporating no bleach activator and (ii) a comparative known bleach activator.

Swatches of 100% cotton 12cm x 12cm were stained using the stains listed below:

	WIL Tea	Tea prepared by Warwick International Group Limited
35	BC1	Tea stains with clay
	BC2	Coffee stains
	BC3	Tea stains
	BC4	Curry stains

## AS4 Chlorophyll stains

These stains are useful to give an indication of the bleaching power of formulations incorporating bleach activator. AS4 is particularly good at indicating  
5 bleaching performance on oily stains.

Wash tests were carried out under conditions typical of both European and U.S. laundry washing conditions. European wash tests were carried out at each of 40°C and 60°C, using cold fill Wascator FOM 71 MP machines,  
10 programmed to BS 4923 (HLCC) wash programmes. U.S. wash tests were carried out at 40°C only, using Maytag (A7500 Model) or Whirlpool machines (Super Capacity Model 3LA5580XS).

For the European wash tests, formulations  
15 incorporating the activator for testing were prepared containing 22.5g PBS4 (sodium perborate tetrahydrate), 124.5g IEC standard base detergent and a weight of activator calculated to give the same moles peracid release as 2% by weight TAED. Formulations were dosed at 7.5g/l  
20 wash water via the dispensing drawer of the machines.

For the U.S. wash tests a wash cycle of 12 minutes duration was used, the temperature of the incoming water being 40°C. Formulations contained 5% by weight activator for testing (100% active) 10% PBS1 (sodium perborate  
25 monohydrate) and IEC to 100%. The formulations were dosed at 1.5g/l.

During the wash tests the swatches were tagged to 2.5kg polyester backing cloth, using plastic tags. After washing the swatches were removed and ironed dry prior to  
30 reading reflectance measurements to assess stain removal. Stain removal was assessed as a percentage brightness and calculated using the equation.

$$\text{Stain removal} = \frac{(RA - RB)}{(RS - RB)} \times 100$$

35

where RA is the reflectance of the washed swatch, RB is the reflectance of the unwashed swatch and RS is the

reflectance of an unstained swatch. Reflectance was measured with U.V. at 420nm using a Spectroflash 500 Spectrophotometer machine.

- 5 Due to variations in swatch make up all evaluations were run compared to a TAED or NBC (N-benzoyl caprolactam) or other conventional activator standard, as well as a blank incorporating no activator.

Table 1

Compound	R Group	R <sup>1</sup> Group	M. Pt (°C)	Purity (HPLC)
A	(CH <sub>2</sub> ) <sub>3</sub>	phenyl	80-91	99.7
B	(CH <sub>2</sub> ) <sub>3</sub>	4-methylphenyl	130-131	99.8
C	(CH <sub>2</sub> ) <sub>3</sub>	4-nitrophenyl	130-131	97.3
D	(CH <sub>2</sub> ) <sub>3</sub>	4-tert-butylphenyl	104.8-105.9	99.4
E	(CH <sub>2</sub> ) <sub>3</sub>	pentyl	liquid	88.1
F	(CH <sub>2</sub> ) <sub>3</sub>	4-chlorophenyl	112.5-114.5	99.3
G	(CH <sub>2</sub> ) <sub>3</sub>	4-methoxyphenyl	119.7-120.1	98.8
H	(CH <sub>2</sub> ) <sub>3</sub>	octyl	liquid	97.3
I	(CH <sub>2</sub> ) <sub>3</sub>	phenoxymethyl	-	99.6
J	(CH <sub>2</sub> ) <sub>3</sub>	3,4-dicarboxyphenyl anhydride	-	-
K	(CH <sub>2</sub> ) <sub>3</sub>	4 tert-butylphenyl	84	100
L	(CH <sub>2</sub> ) <sub>3</sub>	tert-butyl	liquid	95
M	(CH <sub>2</sub> ) <sub>3</sub>	5-(N-phthalimidyl) pentyl	gel	82.9
N	(CH <sub>2</sub> ) <sub>3</sub>	phenoxymethyl	75	99.6
O	(CH <sub>2</sub> ) <sub>6</sub>	phenyl	-	97.4

The results for the wash tests carried out on each of the bleach activators listed in table 1 above, are given in the tables below. In order to ensure that any effect on the results obtained due to variations in water quality could be minimised, for each set of tests run at the same time, a standard and a blank test were also run. The results in the tables below are given in blocks of tests which were carried out at the same time.

The results in tables 2 to 4 below illustrate the effective bleaching properties obtained using the bleach activators of the present invention under the washing conditions tested.

Compound	Wash Test Results (European Washes @ 40°C)					
	WIL TEA	BC1	BC2	BC3	BC4	AS4
Blank	27.9	4.0	11.9	3.7	12.4	31.3
TAED	32.8	9.0	17.4	13.8	12.9	32.0
NBC	37.8	13.4	24.0	18.6	14.5	36.2
A	32.5	10.5	21.7	16.4	16.3	34.6
B	31.7	11.2	22.6	20.2	13.8	39.9
C	38.6	14.6	24.9	24.0	16.3	36.9
Blank	32.0	8.0	14.0	12.9	22.5	22.6
TAED	33.6	10.0	17.8	20.8	21.2	22.3
NBC	41.4	13.1	23.0	28.0	25.3	25.7
D	31.3	10.0	20.2	21.3	22.1	30.7
Blank	29.5	4.9	5.4	8.8	11.6	16.5
TAED	32.6	5.7	7.1	13.9	12.0	17.1
NBC	32.2	9.6	8.7	16.7	14.0	19.1
E	33.6	6.1	7.0	13.0	10.1	19.0
F	36.0	10.7	10.7	20.4	13.6	23.5
Blank	25.7	12.7	12.2	23.2	13.3	19.2
TAED	29.4	12.0	12.9	22.6	13.4	15.5
NBC	31.9	15.2	14.9	24.9	12.6	15.9
G	26.8	15.4	13.8	26.9	14.9	19.5
H	24.1	8.5	10.2	19.8	7.1	17.0
Blank	35.2	4.1	5.1	13.6	10.7	25.1
TAED	40.9	8.4	8.6	21.6	10.3	24.9
NBC	39.6	8.7	8.6	22.6	13.5	27.6
I	32.1	7.4	6.4	20.6	12.2	22.2

TABLE 2 (contd)

Compound	Wash Test Results (European Washes @ 40°C)					
	WIL TEA	BC1	BC2	BC3	BC4	AS4
Blank	30.1	3.6	11.8	5.0	13.6	33.3
TAED	33.6	8.4	17.6	13.8	13.6	32.5
J	38.4	8.0	16.7	9.9	8.4	33.4
Blank	32.0	8.0	14.0	12.9	22.5	22.6
TAED	33.6	10.0	17.8	20.8	21.2	22.3
NBC	41.4	13.1	23.0	28.0	25.3	25.7
K	34.1	9.6	19.1	19.6	20.3	29.8
L	32.0	10.7	19.4	20.3	22.5	24.5
Blank	25.7	12.7	12.2	23.2	13.3	35.1
TAED	29.4	12.0	12.9	22.6	13.4	24.8
NBC	31.9	15.2	14.9	24.9	12.6	15.9
M	22.7	10.7	10.6	18.3	12.2	27.3
Blank	35.2	4.1	5.1	13.6	10.7	25.1
TAED	40.9	8.4	8.6	21.6	10.3	24.9
SNOBS	39.6	8.7	8.6	22.6	13.5	27.6
N	38.6	12.6	11.3	29.9	12.6	26.3
Blank	35.2	4.1	5.1	13.6	10.7	25.1
TAED	40.9	8.4	8.6	21.6	10.3	24.9
NBC	39.6	8.7	8.6	22.6	13.5	27.6
O	38.0	10.2	8.2	23.9	10.3	26.1

TABLE 3

Compound	Wash Test Results (European Washes @ 60°C)					
	WIL TEA	BC1	BC2	BC3	BC4	AS4
Blank	38.6	10.5	18.6	16.1	26.3	44.6
TAED	48.7	18.2	29.4	27.8	25.4	46.4
NBC	44.5	21.2	31.9	31.1	33.2	49.0
A	43.1	17.6	31.7	28.5	31.7	49.7
B	47.3	21.2	36.7	32.2	30.1	52.4
C	53.3	24.5	40.0	37.4	33.1	53.9
Blank	43.0	12.2	23.1	29.5	31.8	32.3
TAED	54.8	18.9	30.8	37.5	34.9	35.0
NBC	54.7	20.3	35.8	41.9	39.6	41.8
D	46.6	18.9	35.2	38.3	42.5	49.6
Blank	40.1	10.7	11.3	23.1	21.3	22.5
TAED	45.4	14.1	14.4	30.4	23.7	23.9
NBC	46.7	15.3	16.7	35.5	24.4	27.1
E	48.2	13.3	14.7	33.2	23.3	25.9
F	46.0	17.2	18.3	38.9	26.2	30.8
Blank	31.9	19.3	15.0	31.7	19.7	29.7
TAED	40.1	21.8	20.2	37.2	20.2	29.4
NBC	41.3	21.4	22.2	38.1	24.8	33.9
G	40.3	22.8	23.7	37.7	26.4	31.7
H	31.9	18.7	17.2	30.0	22.3	30.4
Blank	36.9	11.1	9.8	28.1	22.2	29.9
TAED	43.3	12.4	12.5	34.2	22.0	31.5
NBC	47.7	16.2	16.3	40.1	26.8	34.8
I	40.3	11.8	12.5	32.6	0.6	32.6

TABLE 3 (contd)

Compound	Wash Test Results (European Washes @ 60°C)					
	WIL TEA	BC1	BC2	BC3	BC4	AS4
Blank	38.5	10.6	21.9	19.2	25.1	46.3
TAED	50.3	17.6	31.1	28.8	27.6	44.3
J	49.1	11.4	23.0	19.5	24.1	43.2
Blank	43.0	12.2	23.1	29.5	31.8	32.3
TAED	54.8	18.9	30.8	37.5	34.9	35.0
NBC	54.7	20.3	35.8	41.9	39.6	41.8
K	44.6	19.3	36.5	40.1	41.2	51.5
L	51.5	18.4	32.9	38.5	37.2	38.7
Blank	31.9	19.3	15.0	31.7	19.7	47.4
TAED	40.1	21.8	20.2	37.2	20.2	42.8
SNOBS	41.3	21.4	22.2	38.1	24.8	33.9
M	34.3	17.0	16.2	29.0	19.6	46.4
Blank	36.9	11.1	9.8	28.1	22.2	29.9
TAED	43.3	12.4	12.5	34.2	22.0	31.5
SNOBS	47.7	16.2	16.3	40.1	26.8	34.8
N	n/t	n/t	n/t	n/t	n/t	n/t
Blank	36.9	11.1	9.8	28.1	22.2	29.9
TAED	43.3	12.4	12.5	34.2	22.0	31.5
NBC	47.7	16.2	16.3	40.1	26.8	34.8
O	45.5	15.3	15.3	36.0	25.7	35.0



TABLE 4

Compound	Wash Test Results (US Washes @ 40°C)					
	WIL TEA	BC1	BC2	BC3	BC4	AS4
Blank	10.1	0.2	5.0	0.7	7.7	7.3
TAED	15.5	3.7	6.3	2.4	8.9	8.4
NBC	13.0	4.3	8.8	2.2	8.4	15.3
A	15.0	4.4	8.2	2.2	10.9	13.6
B	9.8	4.3	8.5	3.2	10.2	15.1
C	15.1	4.9	8.9	4.1	11.1	11.5
Blank	10.9	2.3	2.9	-2.3	18.6	5.3
TAED	17.0	2.9	6.3	0.3	15.3	5.9
NBC	11.8	4.6	7.9	-0.5	17.8	5.6
D	7.3	3.9	5.6	-1.8	19.1	8.8
Blank	21.0	2.3	3.1	6.2	8.3	11.0
TAED	28.3	3.8	3.9	9.4	7.6	11.3
SNOBS	23.4	4.3	5.7	11.9	11.1	14.6
E	25.7	4.4	3.8	4.6	7.7	12.2
F	24.0	6.1	4.5	10.3	8.2	13.0
Blank	17.6	5.9	5.9	8.6	6.6	9.1
TAED	23.5	11.3	7.9	9.2	7.8	12.5
SNOBS	18.5	5.5	6.9	9.0	3.4	12.0
G	19.4	10.0	8.8	9.2	6.3	5.0
H	19.0	8.8	5.9	5.8	9.7	9.3
Blank	25.8	1.6	2.4	6.4	7.2	19.4
TAED	32.0	4.5	2.9	7.8	10.2	19.1
SNOBS	29.3	6.1	5.8	10.0	9.2	25.7
I	29.3	3.5	3.3	6.3	11.9	20.0

TABLE 4 (contd)

Compound	Wash Test Results (US Washes @ 40°C)					
	WIL TEA	BC1	BC2	BC3	BC4	AS4
Blank	8.6	0.3	6.9	1.7	11.2	9.9
TAED	17.6	3.4	4.7	2.2	11.4	10.9
SNOBS	17.3	5.0	11.5	1.5	18.8	21.9
J	17.3	2.3	5.2	1.9	9.7	8.4
Blank	10.9	2.3	2.9	-2.3	18.6	5.3
TAED	17.0	2.9	6.3	0.3	15.3	5.9
NBC	11.8	4.6	7.9	-0.5	17.8	5.6
K	6.2	2.5	3.6	-1.5	15.8	7.4
L	5.5	3.2	4.5	-0.7	14.1	6.4
Blank	17.6	5.9	5.9	8.6	6.6	9.1
TAED	23.5	11.3	7.9	9.2	7.8	12.5
SNOBS	18.5	5.5	6.9	9.0	13.4	12.0
M	11.2	6.8	6.9	7.2	3.9	2.7
Blank	25.8	1.6	2.4	6.4	7.2	19.4
TAED	32.0	4.5	2.9	7.8	10.2	19.1
SNOBS	29.3	6.1	5.8	10.0	9.2	25.7
N	26.5	4.5	2.5	6.0	9.1	18.6
Blank	25.8	1.6	2.4	6.4	7.2	19.4
TAED	32.0	4.5	2.9	7.8	10.2	19.1
SNOBS	29.3	6.1	5.8	10.0	9.2	25.7
O	23.8	1.9	-1.1	4.1	7.9	19.4

CLAIMS


1. A bleach activator comprising an optionally substituted N-acyl heterocycle having the formula I:

5




10 in which R represents an optionally substituted alkenyl function having at least 2 carbon atoms in the alkenyl chain which forms a cyclic structure with -N- and -C(O)- and R<sup>1</sup> represents a C<sub>1-30</sub> alkyl or aryl group; or has the formula -R<sup>2</sup>-COX where X may be OH, OM (where M is a metal ion), -ONR<sup>3</sup><sub>4</sub> (where R<sup>3</sup> are each independently selected from H and C<sub>1-4</sub> alkyl), a halogen atom, amine, C<sub>1-30</sub> alkyl or alkoxy or a leaving group and R<sup>2</sup> is a C<sub>1-30</sub> alkylene or arylene group with the proviso that where R is (CH<sub>2</sub>)<sub>5</sub>, R<sup>1</sup> represents a C<sub>3-9</sub> branched alkyl group or a C<sub>10-30</sub> alkyl or aryl group or a C<sub>9-10</sub> branched aralkyl group; or has the formula -R<sup>2</sup>-COX and R<sup>2</sup> is a C<sub>3-10</sub> branched alkyl group or C<sub>11-30</sub> alkylene or arylene group, or a C<sub>9-10</sub> branched alkarylene.

2. A bleach activator according to claim 1 comprising an optionally substituted compound having the formula II:




30 in which R<sup>1</sup> is as defined in claim 1.

3. A bleach activator according to claim 1 comprising an optionally substituted compound of formula III:




35 in which R<sup>1</sup> is as defined in claim 1.

4. A bleach activator according to claim 1 comprising an optionally substituted compound of formula IV:




in which R<sup>1</sup> is as defined in claim 1.

5. A bleach activator according to claim 1 comprising an optionally substituted compound of formula V




in which R<sup>1</sup> is as defined in claim 1.

6. A bleach activator according to claim 1 comprising an optionally substituted compound of formula VI



7. A bleach activator according to claim 1 comprising an optionally substituted compound of formula VII



in which R<sup>1</sup> is as defined in claim 1.


8. A bleach activator according to any preceding claim in which R<sup>1</sup> is an optionally substituted phenyl group, preferably nitrophenyl.

9. A process in which a peroxygen source is reacted with an activator compound according to any preceding claim in a perhydrolysis step in an aqueous solution to form a product solution containing an oxidised product and the product solution is used as a bleach.

10. A composition containing a peroxygen source and an activator according to any of claims 1 to 3.

11. A composition according to claim 10 which is a laundry detergent.

12. A bleaching composition comprising a peroxygen source and at least two bleach activators, at least one of the bleach activators comprising a bleach activator of the formula IX:



in which R represents an alkenyl function having at least 2 carbon atoms in the alkenyl chain which forms a cyclic structure with -N- and -C(O)- and R<sup>1</sup> represents a C<sub>1-30</sub> alkyl or aryl group or has the formula -R<sup>2</sup>-COX where X is OH, OM (where M is a metal ion), -ONR<sup>3</sup>, (where R<sup>3</sup> are each independently selected from H and C<sub>1-4</sub> alkyl), a halogen atom, amine, C<sub>1-30</sub> alkyl or alkoxy or a leaving group as defined for formula I above, and R<sup>2</sup> is a C<sub>1-30</sub> alkylene or arylene group.

10 13. A bleaching composition according to any of claims 10 to 12 comprising a tetra-acylated ethylene diamine, preferably TAED.

14. A bleaching composition according to any of claims 10 to 13 comprising SNOBS.

15

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C11D3/39 C07D227/087

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C11D C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 400 971 (FLORIDA STATE UNIVERSITY) 5 December 1990 see abstract ---	1,2,8
X	WO,A,87 04429 (UPJOHN CO.) 30 July 1987 see page 29, line 24 - line 30; example 1; table 1 ---	1,2
X	EP,A,0 412 058 (SIGMA-TAU IND.) 6 February 1991 see abstract; claim 1 ---	1,3,8
X	EP,A,0 268 218 (NELSON RESEARCH & DEV.) 25 May 1988 see page 5, line 7 - line 39 ---	1,3-5
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

24 March 1995

Date of mailing of the international search report

10. 04. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Grittern, A

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,4 113 735 (V.P. KURKOV) 12 September 1978 see claims ---	1,3
X	DE,A,26 16 993 (BAYER AG) 27 October 1977 see page 13; example 16 ---	1,4,5
A	GB,A,1 596 313 (BOSCH-SIEMENS HAUSGERÄTE GMBH) 26 August 1981 cited in the application see the whole document ---	1,5,9-11
E	WO,A,95 00626 (PROCTER & GAMBLE) 5 January 1995 see claims ---	1,4,8-14
E	WO,A,94 28102 (PROCTER & GAMBLE) 8 December 1994 see claims; example 6 -----	1,5,8-14

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 94/02583

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0400971	05-12-90	US-A- 5175315	29-12-92
		AU-B- 630696	05-11-92
		AU-A- 5515090	20-12-90
		CA-A- 2016951	30-11-90
		CN-A- 1057049	18-12-91
		CN-A- 1086512	11-05-94
		JP-A- 3086860	11-04-91
		NO-C- 174099	16-03-94
		OA-A- 9740	30-11-93
		PL-B- 163946	31-05-94
		US-A- 5336785	09-08-94
WO-A-8704429	30-07-87	AU-A- 6849687	14-08-87
		EP-A- 0232017	12-08-87
		EP-A- 0282492	21-09-88
EP- -0412058	06-02-91	AT-T- 110363	15-09-94
		DE-D- 69011781	29-09-94
		DE-T- 69011781	05-01-95
		ES-T- 2058874	01-11-94
		JP-A- 3024054	01-02-91
		US-A- 5102896	07-04-92
EP-A-0268218	25-05-88	AU-B- 607951	21-03-91
		AU-A- 8132587	26-05-88
		DE-A- 3778833	11-06-92
		US-A- 63150230	22-06-88
US-A-4113735	12-09-78	US-A- 4151171	24-04-79
		US-A- 4151171	24-04-79
		US-A- 4151171	24-04-79
		US-A- 4151171	24-04-79
DE-A-2616993	27-10-77	FR-A- 2348311	10-11-77
		GB-A- 1559627	23-01-80
		JP-A- 52128477	27-10-77
GB-A-1596313	26-08-81	DE-A- 2719235	09-11-78
		FR-A, B 2388924	24-11-78
		SE-B- 439504	17-06-85
		SE-A- 7802908	30-10-78



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 94/02583

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9500626	05-01-95	AU-B- 7061794	17-01-95
WO-A-9428102	08-12-94	AU-B- 6833394	20-12-94

